

# **Dementia in Canada: Prevalence and Incidence 2020 to 2050**

November 2022



**CANADIAN CENTRE FOR  
ECONOMIC ANALYSIS**

## About the Canadian Centre for Economic Analysis

## About the Report

---

The Canadian Centre for Economic Analysis (CANCEA) is a socio-economic research and data firm. CANCEA provides objective, independent and evidence-based analysis and is dedicated to a comprehensive, collaborative, and quantitative understanding of the short- and long-term risks and returns behind market changes, policy decisions and economic behaviour.

CANCEA uses modern techniques in data science, including agent-based modelling, for econometric analysis, risk management assessments, demographic forecasts and epidemiology. CANCEA's work includes market analysis, policy evaluation and risk management, business model optimization, cost-effectiveness and rate of return analysis, macroeconomic analysis, insurance risk evaluation, land use and infrastructure planning, logistics, and labour market analysis. CANCEA also provides comprehensive Canadian data services.

At the centre of CANCEA's analytical capabilities is an agent-based platform called Prosperity at Risk<sup>®</sup> that is an extensive, data-driven model of 56,000 locations across Canada. Given the systems focus behind all of CANCEA's work, CANCEA has a one model approach to its analysis which allows various disciplines and stakeholders to be incorporated into a single analysis.

©2022 Canadian Centre for Economic Analysis  
ISBN: 978-1-989077-39-9  
Printed in Canada • All rights reserved

CANCEA does not accept any research funding or client engagements that require a pre-determined result or policy stance, or otherwise inhibits its independence.

In keeping with CANCEA's guidelines for funded research, the design and method of research, as well as the content of this study, were determined solely by CANCEA.

This information is not intended as specific investment, accounting, legal or tax advice.

**Citation:** Canadian Centre for Economic Analysis, Dementia in Canada: Prevalence and Incidence: 2020 to 2050, 2022

## TABLE OF CONTENTS

Table of Contents .....	i
List of Figures .....	ii
List of Tables .....	ii
Executive Summary.....	1
Prevalence and Incidence of Dementia in Canada .....	1
Informal Caregiving.....	2
Ethnicity and Dementia.....	2
Provincial Differences .....	3
1.0 Introduction .....	5
1.1 Approach.....	5
1.2 About Dementia.....	5
2.0 The Burden of Dementia in Canada, 2020-2050.....	12
2.1 Aging Population .....	12
2.2 Evolving Ethnic Profiles .....	13
2.3 The Burden of Dementia in Canada: 2020-2050.....	15
2.4 Impact of Delaying the Onset of Dementia .....	21
3.0 The Burden of Dementia in Canada's Provinces .....	24
3.1 The Burden of Dementia in Canadian Provinces: 2020-2050 .....	24
3.2 Impact of Delaying the Onset of Dementia .....	27
4.0 Conclusions .....	29
4.1 Data Gaps and Future Research.....	29
A. Methodology.....	31
B. References .....	32

## LIST OF FIGURES

<b>Figure 1</b>	Age standardized incidence rate of dementia (all forms) in Canada.....	7
<b>Figure 2</b>	Age standardized prevalence of dementia (all forms) in Canada.....	8
<b>Figure 3</b>	Population Age Structure: 2020 and 2050.....	12
<b>Figure 4</b>	Ethnic profile of Canadians in 2020 and 2050.....	14
<b>Figure 5</b>	Relative risk of dementia by ethnicity based on (Mayada, et al., 2016).....	14
<b>Figure 6</b>	Prevalence by age and sex, 2020 to 2050.....	16
<b>Figure 7</b>	Annual incidence of any dementia by sex.....	17
<b>Figure 8</b>	Distribution of dementia by ethnic origin by 2050.....	18
<b>Figure 9</b>	Relative change in share of dementia patients from 2020 to 2050 from the analysis.....	18
<b>Figure 10</b>	Number of people with early onset dementia in 2020 and 2050.....	19
<b>Figure 11</b>	Age distribution of informal caregivers, 2020 and 2050.....	20
<b>Figure 12</b>	Impact of delaying the onset of dementia.....	22
<b>Figure 13</b>	Impact of delaying the onset of dementia on the number of informal caregiving hours.....	23
<b>Figure 14</b>	Percentage increase in the number of people with dementia by province; 2020 to 2050.....	25
<b>Figure 15</b>	Total new cases of dementia by type and province from 2020 to 2050.....	25
<b>Figure 16</b>	Change in the number of people with dementia in 2050 relative to 2020.....	27
<b>Figure 17</b>	Reduction in informal caregiving hours in 2050 by province and onset delay.....	28

## LIST OF TABLES

<b>Table 1</b>	The categories and combinations of dementia modelled in the analysis.....	6
<b>Table 2</b>	Potentially modifiable dementia risk factors included in the analysis.....	10
<b>Table 3</b>	Ethnic origins included in the analysis.....	13
<b>Table 4</b>	Prevalence of any dementia by sex, 2020 and 2050.....	16
<b>Table 5</b>	Number of people with dementia, by type, in 2020 and 2050.....	16
<b>Table 6</b>	Total incidence by type, 2020 to 2050.....	17
<b>Table 7</b>	Number of informal caregivers in 2020 and 2050.....	20
<b>Table 8</b>	Number of hours of informal caregivers in 2020 and 2050 (in millions).....	20
<b>Table 9</b>	Number of people with dementia under current trends, and if onset is delayed.....	21
<b>Table 10</b>	Impact of delaying the onset of dementia on the number of informal caregivers.....	22
<b>Table 11</b>	Number of informal caregivers in 2020 and 2050, by province and sex.....	26
<b>Table 12</b>	Annual number of caregiver hours in 2020 and 2050, by province and sex.....	26
<b>Table 13</b>	Reduction in the number of informal caregivers in 2050 compared to current trends.....	28

## EXECUTIVE SUMMARY

The Rising Tide Report, published in 2008, was the first study of its kind to highlight the growing challenge of dementia and to estimate the long-term burden of this disease on the health care and social service systems, on the lives of informal caregivers, and the Canadian economy. Given that over a decade has passed since the seminal work was published, this report revisits the analysis using updated literature estimates of the prevalence and incidence of dementia, along with the most recent relative risk data, for dementia in Canada and its provinces. In addition, the report expands upon the original to take a closer look at the relationship between ethnicity and dementia for both people with dementia and their informal caregivers.

### PREVALENCE AND INCIDENCE OF DEMENTIA IN CANADA

The updated analysis highlights how dementia will continue to be a growing issue in Canada, with the number of people living with some form of dementia expected to almost triple over the next 30 years.

Year	Number of People (65+)				
	Female	Male	Total	Female %	Male %
2020	352,250	217,313	569,563	61.8%	38.2%
2030	592,747	363,167	955,914	62.0%	38.0%
2040	880,677	523,001	1,403,678	62.7%	37.3%
2050	1,053,179	616,039	1,669,218	63.1%	36.9%
Change from 2020-2050	+700,929	+398,725	1,099,655	+1.2%	-1.2%

With the annual number of new diagnoses of dementia expected to more than double by 2050, there could be over 6.3 million new cases of dementia over the next 30 years, with over 60% occurring in women.

Year	Annual New Cases (All Dementia)		
	Female	Male	Total
2020	74,949	48,804	123,753
2030	112,574	74,186	186,760
2040	152,426	98,684	251,110
2050	167,248	109,017	276,265
30 Year Total	3,821,846	2,506,373	6,328,218

However, while there is currently no cure for dementia, many modifiable risk factors can reduce or delay one's risk. To understand the significance of delaying the onset of dementia, three hypothetical scenarios were examined where the onset of dementia was delayed by one, five, or ten years. The analysis highlights how even the modest delay in the onset of dementia of one year could avoid almost one half million new cases of dementia over the next 30 years.

Cumulative New Cases Avoided by 2050				
	Alzheimer's	Other Dementia	Vascular Dementia	Total
Incidence Deferred 1 Year	236,802	138,373	118,857	494,032
Incidence Deferred 5 Years	1,128,449	620,530	538,848	2,287,827
Incidence Deferred 10 Years	2,039,741	1,035,314	955,656	4,030,711

For a larger delay of 10 years, there would be fewer people living with dementia in 2050 than there are today. This drastic shift in prevalence demonstrates that while a cure for dementia might be ideal, any interventions that can delay the onset can also have a significant impact as it would enable people to reach the end of life without developing dementia.

## INFORMAL CAREGIVING

People with dementia are more likely to live in community settings than in long-term care facilities. 69% of people with dementia aged 65 to 80 and 58% aged 80 years and older live outside of long-term care institutions and receive some form of informal caregiving from family or friends. Under these current trends, over 1 million people could be providing informal care to family, relatives, or friends by 2050. This is an increase of 650,000 people relative to 2020. The number of informal caregiving hours could reach almost 1.4 billion hours annually which is equivalent to over 690,000 full-time jobs.

	Year	Current Trends	Incidence Deferred 1 Year	Incidence Deferred 5 Years	Incidence Deferred 10 Years
Number of Informal Caregivers	2020	349,551	344,066	326,251	312,207
	2050	1,005,815	910,569	589,787	307,992
	% Change	188%	165%	81%	-1%
Hours of Informal Care (Millions)	2020	472.6	465.2	441.1	422.1
	2050	1,385.7	1,254.7	813.1	424.9
	% Change	193%	170%	84%	1%

However, this informal caregiving burden would be significantly reduced if the onset of dementia were delayed.


## ETHNICITY AND DEMENTIA

As the ethnic profile of the Canadian population evolves, both through the aging-in place of earlier newcomers and the arrival of recent newcomers it will significantly shift the distribution of incidence for dementia within ethnoculturally diverse populations. In particular, the analysis shows that the representation of Asian immigrants in the population with dementia is expected to triple by 2050, despite their lower risk of acquiring the disorder. In addition, while the total population of people with African origins is a much smaller percentage of the total population, their representation in the population with dementia is expected to more than double. The figure below shows the change in the share of dementia based on the modelling. Values greater than one indicate that their proportion of those with dementia is

## Prevalence and Incidence of Dementia in Canada

larger in 2050 than 2020, while values less than one are the groups whose share is smaller in 2050 than in 2020.

**Ratio of % share of dementia in 2050 and in 2020**



The relationships between ethnicity, dementia, and informal care are complex, and while there is limited literature in the area, as the Canadian population evolves, it is becoming increasingly important to understand and research


### PROVINCIAL DIFFERENCES

---

Across Canada, the growing burden of dementia is not felt uniformly. Several demographic factors combine to result in significant differences in the expected increase in the number of people with dementia by 2050. Provinces with higher growth rates and a faster-aging population (such as Alberta, Ontario, and British Columbia) are expected to see the prevalence of dementia increase faster than the other province in Canada. However, even provinces with slower growth rates, such as the Atlantic Provinces, are expected to see at least a 75% increase in the number of people with dementia.

## Prevalence and Incidence of Dementia in Canada

**Percent growth in number of people with dementia (100% growth means twice as many people)**





## 1.0 INTRODUCTION

Like most developed countries around the world, the Canadian population is undergoing a significant demographic aging process. As the birth rates slow and life expectancy increases, the average age of the population is getting older (Statistics Canada, 2019). Increasing age is associated with an increase risk in developing a variety of age-related diseases and disorders one of which is dementia.

*Rising Tide*, published in 2008, was the first study of its kind to highlight the growing challenge of dementia in Canada, and to estimate the long-term burden of this disease on the health care and social service systems, on the lives of informal caregivers, and on the Canadian economy. Ten years on, new data and research have become available, and more is understood about dementia and its risk factors. This current study aims to incorporate these new research insights, as well as advances in statistical modelling, and recent demographic changes, namely those due to migration, that have occurred over the past decade to generate new projections for the coming decades. Beyond providing a refresh to the 2008 *Rising Tide* report, this study will deepen the analysis with a lens on gender and ethnicity, and focus on informal caregiving.

### 1.1 APPROACH

---

The Landmark Study utilizes CANCEA's socio-economic statistical analysis platform to model and estimate the burden of dementia over the next 30 years in the Canadian population. In the model, each agent is a statistical representation of a person and is associated with several demographic characteristics, including age, sex, ethnicity, which match those of the Canadian population, as informed by publicly available data from Statistics Canada. The initial state of the population is established with age- and sex-specific prevalence of chronic health conditions such as heart disease, smoking, hearing loss, or low-level of physical activity, and the comorbidities between them. The list of risk factors and their prevalence in Canada's population is described in more detail in section 1.2. The model simulates the agents and their interactions over time as they age and pass through various states, such as dementia diagnosis, hospitalization, and death. While similar to other micro-simulation models, such as POHEM (Hennessy, et al., 2015), CANCEA's platform is able to run at the individual level, include interactions and connections between agents (such as family structure) and includes extensive economic accounts. This allows the single model to provide estimates and forecasts of the burden of dementia in Canada over time for measures of health (incidence, prevalence, and mortality) and economics (direct and indirect health care costs, government revenue lost, caregiver costs). This approach enables comparisons of the burden of dementia across different segments of the population, namely by sex and ethnicity, to identify populations particularly vulnerable to dementia in Canada. Additional details are included in Appendix A.

### 1.2 ABOUT DEMENTIA

---

#### 1.2.1 WHAT IS DEMENTIA?

Dementia is a term that encompasses a large class of diseases that are characterized by a decline from a previous cognitive level. Dementia differs from cognitive decline in the course of normal aging in that it

affects day-to-day life and the ability to function in social settings. Symptoms of dementia may include loss of memory, judgment, or reasoning, and changes in mood, behaviour, and ability to communicate. A comprehensive review of dementia can be found in the original Rising Tide report (Smetanin, et al., 2009), Gale et al (2018) or Burns and Iliffe (2009) among others.

An important aspect of dementia is the impact this disease has on those closest to the person affected. Seeing a loved one decline in cognitive ability and experience changes in behaviour can be a difficult experience. In addition, people with dementia can also become more dependent, which can place a high burden of responsibility on family and friends due to the informal caregiving burden, who may also incur significant costs associated with the care of their loved ones.

### 1.2.2 TYPES OF DEMENTIA

There are several different types of dementia. Several reviews on the topic include the original Rising Tide report (Smetanin, et al., 2009), Gale et al (2018) or Burns and Iliffe (2009) among others. These reviews identify Alzheimer's disease which is a progressive degenerative and fatal brain disease in which cell-to-cell connections in the brain are lost, and brain cells eventually die as the most common type of dementia. The second most common type is vascular dementia, which is caused by problems in the supply of blood to the brain due to damage to the vascular system. Conditions that can cause or increase damage to the vascular system include high blood pressure, heart problems, high cholesterol, and diabetes. It is possible to have both Alzheimer's disease and vascular dementia (and combinations with other types of dementia) (Feldman, et al., 2003). Other less common types include dementia with Lewy Bodies, frontotemporal dementia, and Creutzfeldt-Jakob disease. Some dementias occur with chronic non-dementia conditions, such as Parkinson's disease, Huntington's disease. Table 1 shows the categories of dementia included in the analysis, and the percentage of people with the various combinations.

**Table 1** The categories and combinations of dementia modelled in the analysis

Dementia Type	% < 70 years	% > 70 years
Alzheimers Only	47.5%	47.0%
Vascular Dementia Only	9.9%	8.1%
Single Other Dementia	17.9%	6.7%
Alzheimers and Vascular Dementia	9.4%	23.3%
Alzheimers and Other Dementia	7.6%	9.4%
Other combinations	7.7%	5.5%
One or more of any type of dementia	100.0%	100.0%

Derived from Feldman et al (2003)


### 1.2.3 WHO DOES DEMENTIA AFFECT?

Dementia primarily affects people ages 65 and older. However, it can also manifest earlier in life. Globally, there has been an increase in the number of people with dementia, which is associated with increased longevity, and one in three people over 65 now die with dementia (Livingston, et al., 2017). Specific physical health problems (see Section 1.2.5 on Risk Factors) increase the risk of Alzheimer's disease and other forms of dementia. A person with more of these physical illnesses is more likely to develop dementia than those with fewer or no conditions. As a result, many people with dementia also live with chronic co-morbid conditions, and age-related physical health problems (Livingston, et al., 2017).


Dementia is more common in specific segments of the population. Worldwide, there are twice as many women with dementia as men, with women representing 60% of Canada's total people with dementia (Public Health Agency of Canada, 2021). Differences also exist across ethnic and socio-economic groups. Recent studies have shown that Indigenous populations may be at increased risk for dementia compared to non-Indigenous people, the majority population (Jacklin, et al., 2013; Bruce, et al., 2014; Warren, et al., 2015; MacDonald, et al., 2015) and that younger age groups (60 to 79 years old) and males in First Nations communities are disproportionately affected (Jacklin, et al., 2013) These differences can be due in part to the changing population structure within these communities, and Indigenous populations being more affected by the social and environmental factors that increase the risk of developing the disease (Jacklin, et al., 2013; Bruce, et al., 2014; Warren, et al., 2015; MacDonald, et al., 2015).

In Canada, as shown in Figure 1, the age-standardized incidence rate of dementia has been falling since 2009 in both men and women (Public Health Agency of Canada, 2021). As a consequence, the age-standardized prevalence has also fallen slightly, as shown in Figure 2.

**Figure 1** Age standardized incidence rate of dementia (all forms) in Canada (Public Health Agency of Canada, 2021)



**Figure 2** Age standardized prevalence of dementia (all forms) in Canada (Public Health Agency of Canada, 2021)



These most recent data from the Public Health Agency of Canada (PHAC) provide the initial prevalence and incidence of dementia in the analysis for those age 65 and over. Note that provincial age-specific data are used rather than the national aggregates shown in Figure 1 and Figure 2. In addition, the total prevalence of dementia is divided into the categories shown in Table 1. However, dementia can also occur in people under the age of 65. A review by Kurupu & Matthews (2013) reported a prevalence range of 67 to 98 people per 100,000 people in an age group of 45 to 65. Therefore, smoothly extend the incidence and prevalence to younger ages such that they are consistent with both the PHAC and young onset data.

Though there are limited studies investigating the differences in dementia incidence across multiple ethnoracial groups, studies have found evidence of differences in risk profiles (Mayada, et al., 2016). These differences can be complicated by ethnocultural factors, which can influence the perception of the normality of cognitive changes, access to health care, and the level of trust in health care institutions (Babula, et al., 2019). Varying incidence rates between ethnic groups may therefore also be attributable to differences in the timing of diagnosis, clinical presentation and the course of the disease (Babula, et al., 2019).

#### 1.2.4 INFORMAL CAREGIVING

In addition to the direct impact of those with dementia, dementia also has a considerable impact on those close to them. Informal caregiving is care that is provided by family, friends and neighbours as opposed to paid care, which is very common with the majority of people with dementia receiving some form of informal care. Among seniors with dementia, 69% of those younger than 80 and 58% age 80 and older live outside of long-term care homes and receive some informal care (Canadian Institute for Health Information, 2018). Informal caregivers for people with dementia provide an average 26 hours of caregiving per week compared to 17 hours per week for seniors with other health issues (Canadian

Institute for Health Information, 2018). Of those providing informal care, 58% are children and 32% spouses and the remaining 10% are others such as friends or neighbours (Canadian Institute for Health Information, 2018).

However, the likelihood of a person with dementia to receive informal care, and the characteristics of the informal caregiver depends upon many other socio-economic factors beyond age and relationship to the patient. While considerable qualitative research exists using small population samples that focus on specific population groups, or examines caregiving in a broader sense, a quantitative systemic analysis across different ethnoracial groups for informal caregiving for dementia patients in the Canadian context does not exist. In addition, there are other population groups which exhibit differences in caregiving trends. For example, recent research based on the Canadian Longitudinal Study on Aging (Ismail, et al., 2020) showed variations in the likelihood for providing informal care across the LGBTQ communities, with gay and bisexual men more likely to provide informal care than heterosexual men. However, the analysis was not specific to dementia, and additional research would be required to understand better the impact of dementia-specific informal caregiving requirements on the LGBTQ communities.

### 1.2.5 MODIFIABLE RISK FACTORS

The risk factors for developing dementia can be categorized by modifiable and non-modifiable (Smetanin, et al., 2009). Non-modifiable risk factors include aging and genetic risk factors. Potentially modifiable risk factors include physical and mental health conditions, including cardiovascular risk factors, as well as socio-economic, educational, and lifestyle factors (Baumgart, et al., 2015). Updated relative risk values have been reported since the original Rising Tide report. The factors included in the analysis are presented in Table 1 below. Hearing loss has been recently recognized as a risk factor for dementia, with studies showing that even mild levels of hearing loss increase the long-term risk of cognitive decline (Livingston, et al., 2017; Thomson, et al., 2017).

**Table 2** Potentially modifiable dementia risk factors included in the analysis

Risk Factor	Sources	Risk Factor Prevalence
Diabetes	(Gudala, et al., 2013; Ott, et al., 1999; Smolina, et al., 2015; Haroon, et al., 2015)	(Public Health Agency of Canada, 2021)
Obesity	(Tolppanen, et al., 2014)	(Public Health Agency of Canada, 2021)
Stroke	(Kuzma, et al., 2018; Nabalamba & Patten, 2010; Cerasuolo, et al., 2017)	(Public Health Agency of Canada, 2021)
Diet	(Sofi, et al., 2010; Lourida, et al., 2013; Akbaraly, et al., 2019)	(Statistics Canada, 2021)
Hypertension	(Ng, et al., 2013; Tariq & Barber, 2018; Nabalamba & Patten, 2010)	(Public Health Agency of Canada, 2021)
Mood/Anxiety Disorders	(Nabalamba & Patten, 2010; Byers & Yaffe, 2011)	(Public Health Agency of Canada, 2021)
Hearing Loss	(Gurgel, et al., 2014; Thomson, et al., 2017; Lin, et al., 2011; Livingston, et al., 2017)	(Feder, et al., 2015)
Physical Activity	(Scarmeas, et al., 2009; Tariq & Barber, 2018)	(Statistics Canada, 2021)

In addition to the risk factors shown in Table 2, remaining factors such as education and genetics have been incorporated into the base incidence rate. This is equivalent to assuming the the educational profile and genetic risk profile of the population is not changing over the analysis period. Since those at risk of dementia by 2050 would already by born and completed the majority of their education, the relevant genetic and educational profile of population is not changing over time.

### 1.2.6 PREVENTION AND MANAGEMENT

Though common, dementia is not an inevitable part of aging. By some estimates, lifestyle factors could account for one-third of all dementia cases, making these instances theoretically preventable (Livingston, et al., 2017; Baumgart, et al., 2015) While dementia is not presently curable or reversible, it is possible to reduce the risk of developing dementia or delaying its onset, and its symptoms can be managed once diagnosed. As shown in Section 2.4, delaying the onset of dementia even by a few years can represent a substantial gain, as it would enable people to reach the end of life without developing dementia.

Timely diagnosis is the key to managing dementia because it allows for early interventions that can significantly improve the lives of people with dementia and their close ones. Interventions may be pharmacological, psychological, environmental, and social and can help people living with dementia: cope with the aspects of the disorder, disclose the diagnosis to those close to them, maintain or adopt new

## Prevalence and Incidence of Dementia in Canada

daily living activities, handle the cognitive and physical changes and transitions, . In addition, they can help alleviate the burden on caregivers.

## 2.0 THE BURDEN OF DEMENTIA IN CANADA, 2020-2050

Changes in Canada’s demographic landscape are poised to have a significant effect on the dementia prevalence and incidence rates over the next 30 years. A key factor is the change in life expectancy and general aging of the population. Section 2.1 highlights the changes in the population age structure and its implications for growth rates. Another factor is the change in immigration rates and the subsequent change in ethnocultural and visible minority status in Canada. Section 2.2 looks at the evolving diversity in the ethnocultural profile and associated risks for developing dementia. The resulting burden of dementia, in terms of people with dementia, and the impact on informal caregiving for Canada is presented in 2.3 with an examination of provincial differences in Section 3.0. Finally, to further probe the effect of delaying the onset of dementia discussed in Section 0, Section 2.4 presents the results of three hypothetical interventions where the onset of dementia is delayed by 1 year, 5 years, or 10 years.

### 2.1 AGING POPULATION

Since 2000, the median age of Canadians has increased from 36.8 to 40.9 (Statistics Canada, 2019). This trend is expected to continue in the coming decades, and the median age is projected to reach 42.8 by 2050. By that time, the senior population (aged 65+) is estimated to be over 12.5 million, representing an estimated 27% of the population, compared to 18% in 2020. Since age is a risk factor for many chronic health conditions and disabilities, an increase in the number of seniors in the population is expected to lead to an increased prevalence of age-related diseases, and disabilities in the overall population, which could have a significant impact on the Canadian health care and social services systems, and place a disproportionate health and economic burden on informal caregivers.

**Figure 3** Population Age Structure: 2020 and 2050




Figure 3 provides a visual representation of the demographic aging of the Canadian population. Given the link between age and dementia, as the proportion of older citizens in the Canadian population increases,



so will the prevalence and incidence of dementia. In addition, the population of those aged 20 to 50 in 2050, a key demographic group for informal caregiving, is not increasing and is similar to that in 2020.

## 2.2 EVOLVING ETHNIC PROFILES

As discussed in Section 1.2.3 and 1.3.4, ethnicity plays a role in both the incidence of dementia, and the role that informal caregiver performs. Section 2.2.1 discusses the background of the ethnic factors driving the incidence of dementia in the analysis, while Section 2.2.2 discusses the ethnic aspects of informal caregiving.

### 2.2.1 DEMENTIA AND ETHNICITY

The ethnic grouping used in the analysis are taken from the designations of ethnic groupings used by Statistics Canada in the 2016 Population Census (Statistics Canada, 2017). When the census was completed, a long-form question asked participants to self-identify their ethnic origin and year of immigration. This provides the basis for both the ethnic profile of the population and the primary sources of recent immigration. Table 3 presents the ethnic categories used in the analysis.


**Table 3** Ethnic origins included in the analysis

Ethnic Origin	
Oceania origins	North American Aboriginal origins
US American origins	Asian origins
Latin, Central and South American origins	Canadian origins
Caribbean origins	European origins
African origins	

While the primary countries of immigration can change over time, for this analysis, it is assumed that the distribution of countries of origin and age distributions of immigrants will remain similar to that seen over the last decade. Note that while the relative rate of immigration of some ethnic groups, such as those of African origin, has increased compared to historical sources of immigration, such as from Europe, the percentage of the population evolves more slowly. Under these assumptions, Figure 4 shows the evolution of the ethnic profile of Canadians over the next 30 years based on the modelling results. Most notable is the decrease in those with European origins, with a corresponding increase in those of Asian origin.


## Prevalence and Incidence of Dementia in Canada

**Figure 4** Ethnic profile of Canadians in 2020 and 2050



The relative risks of dementia, associated with ethnicity, mapped to the Statistics Canada ethnic groupings relative to those of Asian origins, are shown in Figure 5 (Mayada, et al., 2016). The risks are adjusted to account for other comorbid factors and Mayada found no clear remaining factor to account for the differences between ethnic groups.

**Figure 5** Relative risk of dementia by ethnicity based on (Mayada, et al., 2016)



Of note is the significantly higher risk of dementia for those of African origin. While the percentage of the population with this risk is relatively small, it is growing and dementia rates are expected to increase faster than average in this groups due to the increased risks.

### 2.2.2 INFORMAL CAREGIVING

While large-scale quantitative studies sufficient to estimate reliable relative risks of caregiving based on ethnic origins are limited, there have been many smaller qualitative studies that have examined the various issues that informal caregivers face. A study by Arevalo-Flechas, et al. (2014) found that Latino Americans who provide informal care to people living with Alzheimer's disease experience higher caregiving burden, and cultural values influence, both positively and negatively, how they balance responsibilities. A comparison of African American and White caregivers in the United States found that African-American caregivers provided more hours of care with fewer gender differences (Cohen, et al., 2019), and had differing reasons for providing care (Powers & Whitlatch, 2016). A recent UK study of South Asian caregivers identified several barriers to providing care, including language barriers to services, confusion over research, and feelings of mistrust or stigma (Fry, et al., 2021). A US study revealed that, contrary to expectations about family responsibility in Asian culture, Korean Americans showed favorable attitudes toward using community services in dementia caregiving (Lee & Casado, 2011). As the ethnic profile of Canadians with dementia evolves, these studies highlight the importance of ensuring that obtaining more research on the different needs of these groups to be able to produce culturally safe and sensitive support services and programs to meet these needs.

## 2.3 THE BURDEN OF DEMENTIA IN CANADA: 2020-2050

---

The burden of dementia has two key population groups – those directly affected by the disease and those who provide informal care. Section 2.3.1 examines the prevalence and incidence of dementia from 2020 to 2050, including ethnic profiles and an examination of early-onset dementia. Section 2.3.2 presents the impact of dementia on the provision of informal caregiving including on the the number of people providing informal care, their profiles, and hours of care provided.


### 2.3.1 PREVALENCE AND INCIDENCE

#### 2.3.1.1 OVERALL TRENDS

Over the next 30 years, from 2020 to 2050, the number of people living with dementia in Canada is expected to almost triple from 598,000 today to over 1.7 million in 2050 (Figure 6). While the rate of increase in the number of people with dementia has been slowing compared to a decade ago, the number could still double in 15 years. The number of women with dementia is expected to continue to be significantly greater than the number of men, primarily due to the greater life expectancy of women. In addition, the largest increases in dementia are at the older age groups with the number of people over 85 with dementia expected to increase almost 30% faster than the overall number of cases in the population.

## Prevalence and Incidence of Dementia in Canada

**Figure 6** Prevalence by age and sex, 2020 to 2050



As the Canadian population experiences longer life expectancy, dementia will continue to become more significant. As shown in Table 4, while about 1.6% of the population today has some form of dementia, it is expected to grow to 3.6% by 2050. This growth will put increased pressure on the health care system, long-term care, and informal caregivers.

**Table 4** Prevalence of any dementia by sex, 2020 and 2050

Year	2020		2050		
Gender	Number	% of Population	Number	% of Population	% Growth in Number
Female	366,385	1.9%	1,075,326	4.5%	193%
Male	230,884	1.2%	637,115	2.8%	176%
Total	597,269	1.6%	1,712,441	3.6%	187%

Of the various types of dementia, Alzheimer's will continue to be the dominant with over 1.1 million people by 2050. In comparison, estimates of people living with vascular dementia are approximately 460,000, while 300,00 people are estimated to live with other forms of dementia by 2050. It is important to note that it is possible for an individual to have multiple forms of dementia, so the sum of the subtypes is greater than the population with any form of dementia.

**Table 5** Number of people with dementia, by type, in 2020 and 2050

Year	Alzheimer's	Vascular Dementia	Other Dementia
2020	368,177	156,364	116,828
2050	1,116,594	458,756	300,272
% Increase	203%	193%	157%


Note that a person may have multiple types of dementia so the sum of the individual types will exceed the total with any dementia.

While the number of people living with dementia in 2050 is approximately 1.1 million greater than in 2020, over the next 30 years, there would be over 6.3 million new cases of dementia. The annual number

## Prevalence and Incidence of Dementia in Canada

of new cases could increase from 120,000 new cases each year in Canada today to 276,000 by 2050. This is the equivalent of 329 new cases per day in 2020 and 756 new cases per day by 2050. As the rate of aging of the population slows, the annual number of new cases will start to level off by the 2040s.

**Figure 7** Annual incidence of any dementia by sex



The majority of those are expected to be Alzheimer's, with over 3.2 million diagnoses. In addition, there would be 1.5 million new cases of vascular dementia and 1.6 million new cases of other forms of dementia. Table 6 shows the expected number of new cases of each type of dementia in Canada for each of the next three decades.


**Table 6** Total incidence by type, 2020 to 2050

Years	Alzheimer's	Other Dementia	Vascular Dementia	Total
2020 to 2029	769,011	371,608	354,697	1,495,316
2030 to 2039	1,118,961	533,938	514,380	2,167,279
2040 to 2049	1,360,029	674,954	630,640	2,665,623
<b>Total</b>	<b>3,248,001</b>	<b>1,580,501</b>	<b>1,499,717</b>	<b>6,328,218</b>

### 2.3.1.2 ETHNIC PROFILES


As the population ages, and migration across all ages continues, the ethnic profile of people with dementia will change significantly. These changes arise from historical immigration patterns, future immigration trends of people at risk for dementia, and the varying risk of dementia by ethnicity. Figure 8 shows the expected distribution of people with dementia.

**Figure 8** Distribution of dementia by ethnic origin by 2050



While people of European origin will still be the most common group with dementia by 2050, their proportion of dementia risk is expected to decrease relative to other groups. Conversely as shown in Figure 9, despite having the lowest risk of dementia, the percentage of those with Asian origins living with dementia will triple in the population. This is driven by the largely by the size of the cohort previous immigrants aging in place in this group. Following this trend, the group expected to have the second-highest change in percentage of people living with dementia is of African origin. While the total percentage in the population remains quite small, the elevated risk of developing dementia results in there being twice the prevalence rate in the population.

**Figure 9** Relative change in share of dementia patients from 2020 to 2050 from the analysis. A value of 1 indicates the ethnic group has the same percentage of the dementia cohort in 2050 as in 2020.




The evolving mix of ethnicities of people with dementia will play an important role in determining the requirements and support for those providing informal caregiving and ensuring prompt diagnosis and access to the continuum of care.

### 2.3.1.3 YOUNG ONSET DEMENTIA

While most cases of dementia are in older age groups, by 2050, estimates predicate that there could be over 40,000 people under the age of 65 living with dementia compared to the 28,000 in 2020. These cases represent almost 5% of the current cases of dementia but are not well understood, as the study and identification of early-onset dementia is another area where data and research is underrepresented. The few studies in this area provide very limited estimates of the prevalence and incidence of dementia at younger ages. A review by Kurupu & Matthews (2013) reported a prevalence range of 67 to 98 people per 100,000 people in an age group of 45 to 65 to have a diagnosis of dementia. To better understand the factors driving early onset dementia, and identify any differences in risk factors compared to later onset dementia, additional research is required. This is particularly important when it comes to informal caregiving since many more years of care may be required.

**Figure 10** Number of people with early onset dementia in 2020 and 2050



### 2.3.2 INFORMAL CAREGIVING

People with dementia are likely to have informal caregivers and live outside of long-term care homes in community settings. Among seniors with dementia, 69% of those who are younger than 80, and 58% of those aged 80 and older live outside of long-term care homes (Canadian Institute for Health Information, 2018). Informal caregivers for people with dementia provide an average 26 hours of caregiving per week compared to 17 hours per week for seniors with other health issues (Canadian Institute for Health Information, 2018). Of those providing informal care, 58% are children and 32% spouses, and the remaining 10% are others such as friends or neighbours.

## Prevalence and Incidence of Dementia in Canada


Under the current trends, over 1 million people could be providing informal care to family, relatives, or friends. This is an increase of 650,000 people compared to 349,551 people in 2020.

**Table 7** Number of informal caregivers in 2020 and 2050

	Number of Informal Caregivers			% increase by 2050		
	Male	Female	Both	Male	Female	Both
2020	160,169	189,382	349,551			
2050	467,061	538,754	1,005,815	192%	184%	188%

As shown in Figure 11, there is a large increase in the number of middle-aged people, 45 to 65, highlighting the potential for younger family members to provide care for aging relatives. In addition, there is an increase in older age groups where partners are providing care.

**Figure 11** Age distribution of informal caregivers, 2020 and 2050



Given the large number of people living with dementia, and the level of care required, the number of informal caregiving hours could reach almost 1.4 billion hours annually. This is equivalent to over 690,000 full-time jobs.

**Table 8** Number of hours of informal caregivers in 2020 and 2050 (in millions)

	Number of Caregiving Hours (Millions)			% increase by 2050		
	Male	Female	Both	Male	Female	Both
2020	194.0	278.6	472.6			
2050	579.9	805.8	1,385.7	199%	189%	193%

As shown in the original Rising Tide report, informal caregiving has a sizeable economic impact due to the impact on a caregiver's ability to participate in the workforce.



## 2.4 IMPACT OF DELAYING THE ONSET OF DEMENTIA


While there is no known cure for dementia, there is ongoing research to identify the risk factors and mechanisms that give rise to dementia. In order to understand the significance of even modest improvements in reducing the risks of incidence of dementia and the corresponding delay in disease onset, three hypothetical scenarios were constructed to examine the impact of delaying the onset of dementia by 1, 5 or 10 years. It is important to note that while no specific mechanism is identified to achieve such reductions, the goal is to understand the effects of such interventions. As shown in Table 9, if the onset of dementia is delayed by 10 years, over 4 million new cases of dementia could be avoided by 2050. In this case, there would actually be slightly fewer people living with dementia in 2050 compared to 2020. Even a small delay of 1 year could result in almost 500,000 fewer new cases by 2050.

**Table 9** Number of people with dementia under current trends, and if onset delayed by 1, 5, or 10 years

	Number of People with Any Dementia			% Increase by 2050		
	Male	Female	Both	Male	Female	Both
Current Trends						
2020	230,884	366,385	597,269			
2050	637,115	1,075,326	1,712,441	176%	193%	187%
Incidence Deferred 1 Year						
2020	226,906	360,541	587,447			
2050	573,132	976,891	1,550,023	153%	171%	164%
Incidence Deferred 5 Years						
2020	214,073	341,450	555,523			
2050	359,436	639,705	999,141	68%	87%	80%
Incidence Deferred 10 Years						
2020	204,750	325,669	530,418			
2050	182,049	337,323	519,373	-11%	4%	-2%

## Prevalence and Incidence of Dementia in Canada

**Figure 12** Impact of delaying the onset of dementia




The impact of delaying the onset of dementia on informal caregivers is equally significant. If the onset of dementia were delayed by 10 years, it could reduce number of informal caregiving hours by almost 1 billion hours per year, with 700,000 fewer informal caregivers.

**Table 10** Impact of delaying the onset of dementia on the number of informal caregivers

	Number of Informal Caregivers			% Increase by 2050		
	Male	Female	Both	Male	Female	Both
Current Trends						
2020	160,169	189,382	349,551			
2050	467,061	538,754	1,005,815	192%	184%	188%
Incidence Deferred 1 Year						
2020	157,657	186,409	344,066			
2050	422,878	487,691	910,569	168%	162%	165%
Incidence Deferred 5 Years						
2020	149,509	176,742	326,251			
2050	273,963	315,824	589,787	83%	79%	81%
Incidence Deferred 10 Years						
2020	143,073	169,134	312,207			
2050	143,113	164,879	307,992	0%	-3%	-1%

## Prevalence and Incidence of Dementia in Canada

**Figure 13** Impact of delaying the onset of dementia on the number of informal caregiving hours



## 3.0 THE BURDEN OF DEMENTIA IN CANADA'S PROVINCES

The previous section focused on the burden of dementia at the national level. While there are overall trends that are similar across the provinces, there are also differences unique to each province. Due to their smaller population sizes, the Northwest Territories, Nunavut and the Yukon, with a total population of 126,000 in 2020, are not presented.

### 3.1 THE BURDEN OF DEMENTIA IN CANADIAN PROVINCES: 2020-2050

---

Differences in age distributions, migration patterns, and the prevalence of risk factors combine to produce specific challenges, needs and distribution of burden of dementia for each province. Section 3.1.1 compares the key differences in prevalence and incidence between the provinces, and Section 3.1.2 examines the differences in informal caregiving.

#### 3.1.1 PREVALENCE AND INCIDENCE


As shown in Figure 14, the percentage increase<sup>1</sup> of people living with dementia varies considerably across the country. Faster growing provinces, such as Ontario, Alberta, and British Columbia, have the most significant increases in dementia. In contrast, while the provinces in Atlantic Canada, on average, currently have older populations than many other parts of the country, their much slower population growth limits the rate of growth of dementia. Ontario faces the most new cases with almost 1.5 million new cases of Alzheimer's, 680,000 new cases of vascular dementia, and 780,000 other types of dementia cumulatively by 2050; both Alberta and British Columbia are expected to have a greater percentage increase in people living with dementia.

---

<sup>1</sup> 100% increase corresponds to a doubling in the number of people with dementia


## Prevalence and Incidence of Dementia in Canada

**Figure 14** Percentage increase in the number of people with dementia (any type) by province; 2020 to 2050



Quebec is expected to have the second-highest increase in people with dementia (Figure 15), but stands to be below the national average percentage increase. However, it is important to note that all regions of the country are expected to see an increase in the number of people with dementia if current trends were to continue. Across all provinces, Alzheimer's is the most common form of dementia.

**Figure 15** Total new cases of dementia by type and province from 2020 to 2050



## 3.1.2 INFORMAL CAREGIVING

Across the provinces, the burden of informal care reflects the distribution of people living with dementia. Alberta is expected to see the largest percentage increase in relation to informal caregiving (Table 11) and hours of informal care giving (Table 12) compared to the rest of the country. It is important to note that the number of hours and number of caregivers does not increase by the same amount due to differences in the both the ages of people with dementia, and the characteristics of their caregivers. For example, as people living with dementia age, they may require more hours of informal care. Caregivers who are partners to with the person living with dementia may be able to provide more hours than a child or friend.

**Table 11** Number of informal caregivers in 2020 and 2050, by province and sex

Province	2020			2050			% Increase
	Female	Male	Total	Female	Male	Total	Total
Alberta	15,979	14,337	30,316	62,349	55,801	118,150	290%
BC	23,554	21,992	45,546	74,884	70,046	144,929	218%
Manitoba	5,925	4,289	10,214	13,297	9,655	22,952	125%
NB	3,503	3,424	6,928	6,635	6,509	13,144	90%
NL	1,808	1,511	3,319	3,063	2,662	5,725	72%
Nova Scotia	4,788	4,203	8,991	8,841	7,950	16,792	87%
Ontario	78,186	68,317	146,503	233,762	210,604	444,366	203%
PEI	830	593	1,422	1,978	1,397	3,375	137%
Quebec	49,637	36,469	86,105	121,310	90,277	211,587	146%
Saskatchewan	5,174	5,033	10,207	12,635	12,158	24,794	143%


**Table 12** Annual number of caregiver hours in 2020 and 2050, by province and sex

Province	2020			2050			% Increase
	Female	Male	Total	Female	Male	Total	Total
Alberta	23.2	17.0	40.1	91.7	69.7	161.4	302%
BC	33.8	27.5	61.3	109.6	88.7	198.3	224%
Manitoba	9.1	3.9	13.0	20.8	9.0	29.7	128%
NB	5.6	4.2	9.8	10.8	8.0	18.8	92%
NL	3.5	2.0	5.5	5.9	3.6	9.5	74%
Nova Scotia	7.2	5.3	12.4	13.4	10.1	23.5	90%
Ontario	121.4	89.2	210.6	368.5	276.4	644.9	206%
PEI	1.0	0.4	1.4	2.6	0.9	3.5	142%
Quebec	67.1	39.1	106.1	165.4	100.7	266.1	151%
Saskatchewan	6.9	5.4	12.3	17.0	12.8	29.9	143%

### 3.2 IMPACT OF DELAYING THE ONSET OF DEMENTIA

Across the country, the impact of delaying the onset of dementia would differ for each province. While on average, Canada would see a small decrease in the number of people with dementia by 2050, not all provinces may experience the same effect. In particular, the Atlantic Provinces would experience the largest decreases. Only three provinces, Ontario, British Columbia, and Alberta, would still see increases in the number of people with dementia. As shown in **Error! Reference source not found.**, even small delays in the onset of dementia of one year results in fewer people with dementia across all provinces.

**Figure 16** Change in the number of people with dementia in 2050 relative to 2020 under current trends, and the 3 scenarios where the incidence of dementia is delayed




The reduction in prevalence due to the delay of dementia translates into fewer informal caregivers and caregiving hours. With a 10 year delay, by 2050, Ontario could see 300,000 fewer informal caregivers and 440 million fewer hours of informal caregiving requirements (equivalent to 220,000 full-time jobs). Even a small province like Prince Edward Island would see over 2,000 fewer informal caregivers required in 2050 under the 10 year delay scenario.

## Prevalence and Incidence of Dementia in Canada

**Table 13** Reduction in the number of informal caregivers in 2050 compared to current trends in that year by province and scenario

Province	Incidence Deferred 1 Year	Incidence Deferred 5 Years	Incidence Deferred 10 Years
Alberta	11,460	50,373	83,648
British Columbia	13,503	59,133	99,631
Manitoba	2,333	9,809	16,261
New Brunswick	1,273	5,635	9,313
Newfoundland and Labrador	565	2,439	3,982
Nova Scotia	1,621	7,151	11,842
Ontario	41,067	180,959	305,396
Prince Edward Island	330	1,366	2,308
Quebec	20,759	88,940	148,269
Saskatchewan	2,334	10,223	17,173
Canada	95,246	416,028	697,823

**Figure 17** Reduction in informal caregiving hours in 2050 by province and onset delay





## 4.0 CONCLUSIONS

Dementia will continue to be a growing issue in Canada, with the number of people living with some form of dementia almost tripling over the next 30 years. Under the current trends, 1.7 million people could be living with dementia by 2050, with over 6.3 million new diagnoses of dementia occurring from 2020 to 2050. While over 51% of the new cases are expected to be Alzheimer's, vascular dementia would constitute almost 24% of the new cases, with other forms of dementia making up the remainder.

Over this period, the ethnic profile of people with dementia could change significantly. This changing ethnic profile may require different approaches to identify, diagnose, and manage dementia due to the variety of cultural, social, and economic factors that impact people living with dementia, and informal caregivers. By 2050, over 650,000 additional people (relative to 2020) would be providing informal care and totalling almost 1.4 billion hours annually. This informal caregiving burden is equivalent to 690,000 full-time jobs.

The growing informal caregiving requirements have a significant burden on families, relatives, and friends, not only due to the hours required, but also impacting many people in their key working years, this can have an effect on personal finances and overall economic activity. .

While all provinces of the country are expected to see an increase in the burden of dementia if current trends continue, the distribution of the burden of dementia will not be uniform. Provinces with higher growth rates and a faster aging population (such as Alberta, Ontario, and British Columbia) are expected to see the prevalence of dementia increase faster than the other provinces in Canada. These regional differences highlight the risk of assuming that dementia is evolving the same uniformly across the country and the need to ensure any health or caregiving interventions are appropriately targeted to each province.

Any interventions which result even in small delays in the onset of dementia could significantly reduce the number of people living with dementia, and the informal caregiving burden. If the onset of dementia is delayed by 1 year, it would avoid almost 500,000 cases of dementia over the next 30 years. If the incidence of dementia were delayed by 10 years, over 4 million cases of dementia could be avoided and most regions of the country would see fewer people living with dementia in 2050 than today.

### 4.1 DATA GAPS AND FUTURE RESEARCH

---

As with many quantitative modelling analysis projects, the process of parameterizing the model brings to light areas with data gaps. While the disease dynamics related to dementia are well studied in terms of risk factors, prevalence and incidence in older age groups, the same level of detail is not available for younger people with dementia. Given that these younger people with dementia age into the older cohorts, an understanding of the incidence rates and risks for early onset dementia is important to better quantify the future burden of dementia, and in particular the effects of any early interventions.

In addition, more nuanced data on the many aspects of informal caregiving is required to better understand the relationships between people with dementia and their caregivers. In particular, quantitative data on any ethnic differences of caregiving is lacking, such as the relative likelihood of

## Prevalence and Incidence of Dementia in Canada

informal caregiving across a range of ethnic backgrounds. While qualitative studies indicate that factors such as the hours of informal caregiving, or the relationship between caregiver and dementia patient, may vary among ethnic groups, there is a lack of comprehensive comparative data across a variety of ethnicities. Beyond ethnic groups, other population groups and socio-economic groups, such as those in the LGBTQ2I communities, may have different experiences and requirements for providing informal care for people living with dementia, but quantitative data is lacking.

As the demographics of people living with dementia, and r caregivers change, ongoing research will be critical to understanding how it will affect both the health care system and those providing informal care in the future.

## A. METHODOLOGY

The analysis was conducted using CANCEA's agent-based data analysis and modelling platform. In the platform, every individual in the country is tracked over 30 years with every diseases and risk factor modelled for each person. Over the course of an individual's simulated life, they have an evolving likelihood of getting dementia, or changing any other health state, depending on age, sex, and the other health states for the individual. In addition to the evolution of the initial population, people exit the simulation through either death or emigration. Birth and immigration add new people to the model. The conditional probabilities of each change of state is determined through training (machine learning) the platform on historical data so that history can be reproduced.

A complete description of the methodology can be found at <https://www.cancea.ca>.

## B. REFERENCES

- Akbaraly, T. et al., 2019. Association of Midlife Diet With Subsequent Risk for Dementia. *JAMA*, 321(10), pp. 957-968.
- Arevalo-Flechas, L. et al., 2014. Latino Alzheimer's caregivers: what is important to them?. *Journal of Managerial Psychology*, 29(6), pp. 661-684.
- Babula, G. et al., 2019. Perspectives on ethnic and racial disparities in Alzheimer's disease and related dementias: Update and areas of immediate need. *Alzheimer's & Dementia*, 15(2), pp. 292-312.
- Baumgart, M. et al., 2015. Summary of the evidence on modifiable risk factors for cognitive decline and dementia: A population-based perspective. *Alzheimer's & Dementia*, 11(6), pp. 718-726.
- Bruce, S., Riediger, N. & Lix, L., 2014. Chronic disease and chronic disease risk factors among First Nations, Inuit and Metis populations of northern Canada. *Chronic Diseases and Injuries in Canada*, 34(4), pp. 210-217.
- Burns, A. & Iliffe, S., 2009. Dementia. *BMJ*, 338(b75), pp. 405-409.
- Byers, A. & Yaffe, K., 2011. Depression and risk of developing dementia. *Nature Reviews Neurology*, Volume 7, pp. 323-31.
- Canadian Institute for Health Information, 2018. *Dementia in Canada*, Ottawa: CIHI.
- Cerasuolo, J. et al., 2017. Population-based stroke and dementia incidence trends: Age and sex variations. *Alzheimer's Dement.*, 13(10), pp. 1081-88.
- Cohen, S. et al., 2019. Differences within Differences: Gender Inequalities in Caregiving Intensity Vary by Race and Ethnicity in Informal Caregivers. *Journal of Cross-Cultural Gerontology*, Volume 34, pp. 245-263.
- Feder, K. et al., 2015. Prevalence of hearing loss among Canadians aged 20 to 79: Audiometric results from the 2012/2013 Canadian Health Measures Survey. *Health Rep.*, 26(7), pp. 18-25.
- Feldman, H. et al., 2003. A Canadian cohort study of cognitive impairment and related dementias (ACCORD): study methods and baseline results. *Neuroepidemiology*, 22(5), pp. 265-74.
- Fry, G. et al., 2021. Recruitment of south Asian carers into a survey-based research study. *Dementia (London)*, 20(3), pp. 1154-1161.
- Gale, S., Acar, D. & Daffner, K., 2018. Dementia. *The American Journal of Medicine*, 131(10), pp. 1161-1169.
- Gudala, K., Bansal, D., Schifano, F. & Bhansali, A., 2013. Diabetes mellitus and risk of dementia: A meta-analysis of prospective observational studies. *J Diabetes Investig*, 4(6), pp. 640-650.

Gurgel, R. et al., 2014. Relationship of Hearing loss and Dementia: a Prospective, Population-based Study. *Otol Neurotol.*, 35(5), pp. 775-781.

Haroon, N. et al., 2015. Risk of dementia in seniors with newly diagnosed diabetes: a population-based study. *Diabetes Care*, 38(10), pp. 1868-75.

Hennessy, D. et al., 2015. The Population Health Model (POHEM): an overview of rationale, methods and applications. *Population Health Metrics*, 13(24).

Ismail, M., Hammond, N., Wilson, K. & Stinchombe, A., 2020. Canadians Who Care: Social Networks and Informal Caregiving Among Lesbian, Gay, and Bisexual Older Adults in the Canadian Longitudinal Study on Aging. *The International Journal of Aging and Human Development*, 91(3), pp. 299-316.

Jacklin, K., Walker, J. & Shawande, M., 2013. The Emergence of Dementia as a Health Concern Among First Nations Populations in Alberta, Canada. *Can J Public Health*, 104(1), pp. e39-e44.

Kurupu, D. & Matthews, B., 2013. Young-Onset Dementia. *Semin Neurol.*, 33(4), pp. 365-385.

Kuzma, E. et al., 2018. Stroke and dementia risk: A systematic review and meta-analysis. *Alzheimers Dement.*, 14(11), pp. 1416-26.

Lee, S. & Casado, B., 2011. Attitudes toward Community Services Use in Dementia Care among Korean Americans. *Clinical Gerontologist*, 34(4), pp. 271-286.

Lin, F. et al., 2011. Hearing Loss and Incident Dementia. *Arch Neurol*, 68(2), pp. 214-220.

Livingston, G. et al., 2017. Dementia prevention, intervention, and care. *Lancet*, Volume 390, pp. 2673-734.

Lourida, I. et al., 2013. Mediterranean Diet, Cognitive Function, and Dementia: A Systematic Review. *Epidemiology*, 24(4), pp. 479-489.

MacDonald, J., Barnes, D. & Middleton, L., 2015. Implications of Risk Factors for Alzheimer's Disease in Canada's Indigenous Population. *Can Geriatr J*, 18(3), pp. 152-158.

Mayada, E., Glymour, M. M., Quesenberry, C. & Whitmer, R., 2016. Inequalities in dementia incidence between six racial and ethnic groups over 14 years. *Alzheimers Dement.*, 12(3), pp. 216-224.

Nabalamba, A. & Patten, S., 2010. Prevalence of mental disorders in a Canadian household population with dementia. *Can J Neurol Sci*, 37(2), pp. 186-94.

Ng, J., Turek, M. & Hakim, A., 2013. Heart disease as a risk factor for dementia. *Clinical Epidemiology*, Volume 5, pp. 135-145.

Ott, A. et al., 1999. Diabetes mellitus and the risk of dementia: The Rotterdam Study. *Neurology*, 53(9), pp. 1937-42.

Powers, S. & Whitlatch, C., 2016. Measuring cultural justifications for caregiving in African American and White caregivers. *Dementia (London)*, 15(4), pp. 629-45.

Public Health Agency of Canada, 2021. *Canadian Chronic Disease Surveillance System*. [Online] Available at: <https://health-infobase.canada.ca/ccdss/data-tool/> [Accessed 01 09 2021].

Scarmeas, N. et al., 2009. Physical activity, diet, and risk of Alzheimer disease. *JAMA*, 302(6), pp. 627-37.

Smetanin, P. et al., 2009. *Rising Tide: The Impact of Dementia in Canada 2008 to 2038*, Toronto: RiskAnalytica.

Smolina, K., Wotton, C. & Goldacre, M., 2015. Risk of dementia in patients hospitalised with type 1 and type 2 diabetes in England, 1998-2011: a retrospective national record linkage cohort study. *Diabetologia*, 58(5), pp. 942-50.

Sofi, F., Abbate, R., Gensini, G. F. & Casini, A., 2010. Accruing evidence on benefits of adherence to the Mediterranean diet on health: an updated systematic review and meta-analysis. *Am J Clin Nutr*, 92(5), pp. 1189-96.

Statistics Canada, 2017. *Census Profile. 2016 Census*, s.l.: Statistics Canada.

Statistics Canada, 2019. *Population Projections for Canada (2018 to 2068), Provinces and Territories (2018 to 2043)*, Ottawa: s.n.

Statistics Canada, 2021. *Table 13-10-0096-01 Health characteristics, annual estimates*, s.l.: s.n.

Tariq, S. & Barber, P., 2018. Dementia risk and prevention by targeting modifiable vascular risk factors. *J Neurochem*, 144(5), pp. 565-81.

Thomson, R., Auduong, P., Miller, A. & Gurgel, R., 2017. Hearing loss as a risk factor for dementia: A systematic review. *Laryngoscope Investig Otolaryngol*, 2(2), pp. 69-79.

Tolppanen, A.-M. et al., 2014. Midlife and late-life body mass index and late-life dementia: results from a prospective population-based cohort. *J Alzheimers Dis*, 38(1), pp. 201-9.

Warren, L. et al., 2015. Prevalence and incidence of dementia among indigenous populations: a systematic review. *International Psychogeriatrics*, 27(12), pp. 1-12.